

ON THE ÉTALE FUNDAMENTAL GROUPS OF ARITHMETIC SCHEMES

FENG-WEN AN

ABSTRACT. In this paper we will give a computation of the étale fundamental group of an integral arithmetic scheme. For such a scheme, we will prove that the étale fundamental group is naturally isomorphic to the Galois group of the maximal formally unramified extension over the function field. It consists of the main theorem of the paper. Here, formally unramified will be proved to be arithmetically unramified which is defined in an evident manner and coincides with that in algebraic number theory. Hence, formally unramified has an arithmetic sense. At the same time, such a computation coincides with the known result for a normal noetherian scheme.

Contents

Introduction	
1. Statement of The Main Theorem	
2. Preliminaries	
3. A Criterion of Galois Covers	
4. The Calculus and Construction of Galois Covers	
5. Formally Unramified = Arithmetically Unramified	
6. Proof of The Main Theorem	
References	

INTRODUCTION

In this paper we will give a computation of the étale fundamental group of an arithmetic variety, i.e., an integral arithmetic scheme surjectively over $\text{Spec}(\mathbb{Z})$ of finite type.

Let X be an arithmetic variety. It will be proved that the étale fundamental group $\pi_1^{et}(X, s)$ is isomorphic to the Galois group $\text{Gal}(k(X)^{un}/k(X))$ of the maximal formally unramified extension $k(X)^{un}$ of the function field $k(X)$ in a natural manner. See the main theorem of the paper.

Such a computation of the étale fundamental group coincides with the known result on a normal noetherian scheme (for instance, see [9, 11, 15, 18]).

In particular, we will prove that the maximal formally unramified extension $k(X)^{un}$ is equal to the maximal arithmetically unramified extension $k(X)^{au}$.

Here, arithmetically unramified, which is defined in an evident manner (see §5), coincides with that in algebraic number theory and has many arithmetic operations and properties. Hence, the maximal formally unramified extension is exactly of an arithmetic sense.

2010 *Mathematics Subject Classification.* Primary 14F35; Secondary 11G35.

Key words and phrases. arithmetic scheme, arithmetically unramified, étale fundamental group, formally unramified.

At the same time, the arithmetically unramified is more accessible to operations. Naturally, the arithmetically unramified can serve as a computation of the étale fundamental group of an arithmetic variety.

For instance, as an immediate application, take a finite number of variables $t_1, t_2, \dots, t_n$ over $\mathbb{Q}$. It is proved that $\mathbb{Q}(t_1, t_2, \dots, t_n)$ has a trivial maximal arithmetically unramified extension (see [7]). Therefore, the arithmetic variety $\text{Spec}(\mathbb{Z}[t_1, t_2, \dots, t_n])$ has a trivial étale fundamental group (see *Example 1.8*), i.e.,

$$\pi_1^{et}(\text{Spec}(\mathbb{Z}[t_1, t_2, \dots, t_n])) \cong \{0\}.$$

1. STATEMENT OF THE MAIN THEOREM

1.1. Notations. In this paper, an **arithmetic variety** is an integral scheme that is surjectively over $\text{Spec}(\mathbb{Z})$ of finite type.

For an integral scheme Y , let $k(Y) = \mathcal{O}_{Y, \xi}$ be the function field of Y , where ξ is the generic point of Y . Denote by $\pi_1^{et}(Y, s)$ the étale fundamental group of Y over a geometric point s of Y over a given algebraic closure of the function field $k(Y)$. Sometimes, we write $\pi_1^{et}(Y) = \pi_1^{et}(Y, s)$ for brevity.

Let $\text{Gal}(L/K)$ denote the Galois group of an extension L of a field K .

1.2. Formally unramified. Fixed an arithmetic variety X . Let Ω be an algebraic closure of the function field $k(X)$ and s_ξ the geometric point of X over Ω such that $s_\xi(\text{Spec}(\Omega))$ is the generic point ξ of X .

Put

- $[X; \Omega]_{et} \triangleq$ the set of the equivalence classes of pointed covers (Y, s_Y) of (X, s_ξ) such that Y is an irreducible finite étale Galois cover of X and s_Y is a geometric point of Y over s_ξ . Here, two such covers $(Y, s_Y), (Z, s_Z)$ of (X, s_ξ) are equivalent if there is an isomorphism f of Y onto Z over X with $f(s_Y) = s_Z$.

For finite étale Galois covers, see §6.1 below in the paper or see [9, 11, 15, 18].

For brevity, denote by Y or (Y, s_Y) for an equivalence class of the pointed cover (Y, s_Y) that is contained in $[X; \Omega]_{et}$.

For any $X_\alpha, X_\beta \in [X; \Omega]_{et}$, we say

$$X_\alpha \leq X_\beta$$

if and only if X_β is a finite étale Galois cover over X_α . Then $[X; \Omega]_{et}$ is a directed set with the partial order $\leq$.

Set

- $k(X; \Omega)^{un} \triangleq \lim_{\rightarrow Z \in [X; \Omega]_{et}} \lambda_Z(k(Z))$, i.e., $k(X; \Omega)^{un}$ is the direct limit of the direct system of fields $k(Z)$ indexed by $[X; \Omega]_{et}$, where each $\lambda_Z : k(Z) \rightarrow \Omega$ is the $k(X)$ -embedding of fields.

The subfield $k(X; \Omega)^{un}$ of Ω is said to be the **maximal formally unramified extension** of the arithmetic variety X (or of the function field $k(X)$) in Ω .

It is seen that $k(X; \Omega)^{un}$ is well-defined since all the $k(X)$ -embeddings

$$\lambda_Z : k(Z) \rightarrow \Omega$$

are compatible in a natural manner. Moreover, $k(X; \Omega)^{un}$ is an algebraic Galois extension of the field $k(X)$.

In deed, it will also be seen that the maximal formally unramified extension is exactly of an arithmetic sense that coincide with algebraic number theory (see §§1.4, 5.2 for details; see [7, 8] for further results).

1.3. Statement of the main theorem. Now we give a computation of the étale fundamental group of an arithmetic variety, which is the main theorem of the paper.

Theorem 1.1. *For an arithmetic variety X , the étale fundamental group $\pi_1^{et}(X, s)$ is naturally isomorphic to the Galois group of the maximal arithmetically unramified extension $k(X; \Omega)^{un}$ over the function field $k(X)$.*

That is, there is an isomorphism

$$\pi_1^{et}(X, s) \cong \text{Gal}(k(X; \Omega)^{un}/k(X))$$

between groups in a natural manner.

Here, s is a geometric point of X over an algebraic closure Ω of the function field $k(X)$.

Remark 1.2. Such a computation above coincides with the known results on étale fundamental groups of normal connected schemes (e.g., see [5, 9, 11, 15, 18]).

We will prove *Theorem 1.1* in §6 after we make preparations in §§2-5.

1.4. Concluding Remarks. There are the following remarks as a conclusion of the section.

Remark 1.3. The étale fundamental groups of arithmetic varieties are a birational invariant. In deed, let X and Y be two arithmetic varieties such that the function fields $k(X) \cong k(Y)$ are isomorphic. Then there is a natural isomorphism

$$\pi_1^{et}(X) \cong \pi_1^{et}(Y)$$

between the étale fundamental groups.

Remark 1.4. Fixed an arithmetic variety X and an algebraic closure Ω of $k(X)$. Let $k(X, \Omega)^{au}$ be the the maximal arithmetically unramified extension of $k(X)$ in Ω (see §5 for detail). It is seen that

$$k(X; \Omega)^{au} = k(X; \Omega)^{un}$$

holds, i.e., for the field $k(X)$, the maximal formally unramified extension is equal to the maximal arithmetically unramified extension that is given in an arithmetic manner (see *Lemma 5.9*). Hence, the formally unramified extension $k(X; \Omega)^{un}$ has a sense of arithmetics.

Remark 1.5. In [3] we also give a computation of the étale fundamental group of an algebraic scheme by virtue of algebraic unramified extensions over the function field. In deed, let Z be an integral algebraic scheme Z over a field K of finite type. It is seen that the étale fundamental group $\pi_1^{et}(Z, s)$ is naturally isomorphic to the Galois group of the maximal algebraically unramified extension over the function field $k(Z)$. However, the algebraically unramified and the arithmetically unramified are very different from each other. In particular, the algebraically unramified is different from that in algebraic number theory.

Remark 1.6. Let X be an arithmetic variety and let $\mathcal{O}_{k(X)}$ be the ring of algebraic integers in the field $k(X)$ (see §5 for detail). There is an isomorphism

$$\pi_1^{et}(X) \cong \pi_1^{et}(\text{Spec}(\mathcal{O}_{k(X)}))$$

in a natural manner. Hence, to compute $\pi_1^{et}(X)$, canonically it reduces to find the étale fundamental group of the affine scheme $\text{Spec}(\mathcal{O}_{k(X)})$.

Remark 1.7. Let X be an arithmetic variety. Put $L = k(X)$. Then there are isomorphisms

$$\pi_1^{et}(X) \cong \text{Gal}(L^{au}/L) \cong \text{Gal}(L^{un}/L)$$

of groups in canonical manner from *Theorem 1.1*, *Remarks 1.4*, *5.10*.

At the end of the section, as an application of the main theorem in the paper, we give a computation of the étale fundamental group of the ring of algebraic integers in a purely transcendental extension over $\mathbb{Q}$.

Example 1.8. Let $t_1, t_2, \dots, t_n$ be variables over $\mathbb{Q}$. Then

$$\pi_1^{et}(\text{Spec}(\mathbb{Z}[t_1, t_2, \dots, t_n])) \cong \{0\}$$

since

$$\mathbb{Q}(t_1, t_2, \dots, t_n)^{au} = \mathbb{Q}(t_1, t_2, \dots, t_n)$$

holds (see [7]).

2. PRELIMINARIES

In the following we will fix notations and terminology which will be used to prove the main theorem of the paper.

2.1. Notations. In the remainder of the paper, by an **integral variety** we will understand a finite-dimensional integral scheme that is surjectively over $\text{Spec}(\mathbb{Z})$.

Define L^{al} (or $\overline{L}$) $\triangleq$ an algebraical closure of a field L ; $Fr(D)$ $\triangleq$ the field of fractions of an integral domain D .

Let $f : X \rightarrow Y$ be a morphism of integral schemes. Denote by $f^\# : k(Y) \rightarrow k(X)$ the homomorphism of the function fields given by f ; put $\text{Aut}(X/Y)$ $\triangleq$ the group of automorphisms of X over Y .

Let A be a commutative ring. Set

- $j_x(A)$ (or j_x) $\triangleq$ the prime ideal of A corresponding to a point $x \in \text{Spec}(A)$.

2.2. Essentially affine schemes. Let X be a scheme. An **affine covering** of X is a family

$$\mathcal{C}_X = \{(U_\alpha, \phi_\alpha; A_\alpha)\}_{\alpha \in \Delta}$$

where for each $\alpha \in \Delta$, ϕ_α is an isomorphism from an open set U_α of X onto the spectrum $\text{Spec } A_\alpha$ of a commutative ring A_α . Each $(U_\alpha, \phi_\alpha; A_\alpha) \in \mathcal{C}_X$ is called a **local chart**. For brevity, a local chart $(U_\alpha, \phi_\alpha; A_\alpha)$ will sometimes be denoted by U_α or (U_α, ϕ_α) .

An affine covering $\mathcal{C}_X$ of $(X, \mathcal{O}_X)$ is said to be **reduced** if $U_\alpha \neq U_\beta$ holds for any $\alpha \neq \beta$ in Δ .

Let **Comm** be the category of commutative rings with identity. For a given field Ω , let **Comm**(Ω) be the category consisting of the subrings of Ω and their isomorphisms.

Definition 2.1. Let $\mathfrak{Comm}_0$ be a subcategory of $\mathfrak{Comm}$. An affine covering of X , namely $\{(U_\alpha, \phi_\alpha; A_\alpha)\}_{\alpha \in \Delta}$, is said to be **with values** in $\mathfrak{Comm}_0$ if for each $\alpha \in \Delta$, there are the properties:

$$\mathcal{O}_X(U_\alpha) = A_\alpha;$$

$$U_\alpha = \text{Spec}(A_\alpha),$$

where each A_α is a ring contained in $\mathfrak{Comm}_0$.

In particular, an affine covering $\mathcal{C}_X$ of X with values in $\mathfrak{Comm}(\Omega)$ is said to be **with values in the field Ω** .

Definition 2.2. An integral scheme X is said to be **essentially affine** in a field Ω if for every affine open set U of X there are the properties:

$$\mathcal{O}_X(U) \subseteq \Omega;$$

$$U = \text{Spec}(\mathcal{O}_X(U)).$$

Here we give such a fact that every integral variety X is isomorphic to an integral variety Z that is essentially affine in the field $k(X)$.

Lemma 2.3. *For any integral variety X , there is an integral variety Z satisfying the properties:*

- $k(X) = k(Z)$;
- $X \cong Z$ are isomorphic schemes;
- Z is essentially affine in the field $k(Z)$.

Such an integral variety Z is called an **essentially affine realization** of X .

Proof. Fixed an affine open set U of Y . Let

$$\phi_U : (U, \mathcal{O}_X|_U) \rightarrow (\text{Spec}(A_U), \mathcal{O}_{\text{Spec}(A_U)})$$

be an isomorphism of schemes. We have ring isomorphisms

$$\phi_U^\# : A_U \rightarrow \mathcal{O}_X(U);$$

$$i_X : \mathcal{O}_X(U) \rightarrow B_U \subseteq k(X).$$

The composite of ring isomorphisms

$$t_U = i_X \circ \phi_U^\# : A_U \rightarrow B_U$$

induces an isomorphism

$$\tau_U : (\text{Spec}(A_U), \mathcal{O}_{\text{Spec}(A_U)}) \rightarrow (\text{Spec}(B_U), \mathcal{O}_{\text{Spec}(B_U)})$$

between affine schemes. Put

$$\psi_U = \tau_U \circ \phi_U.$$

Then there is a new scheme Z obtained by gluing all such affine schemes

$$(\text{Spec}(B_U), \mathcal{O}_{\text{Spec}(B_U)})$$

along ψ_U for every affine open set U of X (see [10, 12, 13]). It is seen that the scheme Z has the desired properties. $\square$

2.3. Affine points in a field. The affine points in a field will be applied to a universal construction of a Galois cover of an integral variety in §4. Let Ψ be an arbitrary field.

Definition 2.4. By an **affine Ψ -point** (or **affine point** in Ψ) we understand a point contained in the underlying space of an affine scheme $\text{Spec}(A)$ for a subring A of the field Ψ such that $\text{Fr}(A) = \Psi$.

Denote by $[\Psi]_{ap}$ the set of all affine points in the field Ψ .

In general, it is not true that there is an affine scheme that has $[\Psi]_{ap}$ as the underlying space.

Definition 2.5. Let $x, y \in [\Psi]_{ap}$ be two affine points.

(i) x and y are **Ψ -equal** if there is a subring A of the field Ψ such that

- $\text{Fr}(A) = \Psi$;
- $x, y \in \text{Spec}(A)$;
- $j_x(A) = j_y(A)$ holds as prime ideals in A .

(ii) x and y are **Ψ -equivalent** if there are a tower of subrings $A \subseteq B \subseteq \Psi$ satisfying the properties:

- $\text{Fr}(A) = \text{Fr}(B) = \Psi$;
- $x \in \text{Spec}(A)$, $y \in \text{Spec}(B)$;
- $j_x(A) = \rho_{A,B}^{-1}(j_y(B))$ holds as prime ideals in A ,

where $\rho_{A,B} : A \hookrightarrow B$ is the inclusion map.

It is clear that affine Ψ -points that are Ψ -equal are Ψ -equivalent.

2.4. Function fields and invariant subrings. Let X be an integral variety with generic point ξ . Recall that for the function field $k(X)$, there is a natural inclusion

$$i_X : \mathcal{O}_X(U) \rightarrow k(X) \triangleq \mathcal{O}_{X,\xi}$$

of rings for each open set U in X , i.e.,

$$k(X) = \{i_X(x) \in k(X) : x \in \mathcal{O}_X(U), U \text{ is open in } X\}.$$

Furthermore, $k(X)$ is isomorphic onto the field

$$\text{Rat}(X) \triangleq \{[U, f] : f \in \mathcal{O}_X(U), U \text{ is open in } X\},$$

where $[U, f]$ denotes the germ of (U, f) (see [10, 12, 13]).

For brevity, we will always identify $k(X)$, (U, f) , and $i_X(x) \in k(X)$ with $\text{Rat}(X)$, $[U, f]$, and $x \in \mathcal{O}_X(U)$, respectively.

Each automorphism $\sigma = (\sigma, \sigma^\#)$ of X gives an automorphism

$$\sigma^\# \triangleq \sigma_\xi^\# : \mathcal{O}_{X,\xi} \rightarrow \mathcal{O}_{X,\xi}$$

of the function field $k(X)$.

For an open set U in X and a subgroup $G \subseteq \text{Aut}(X/Y)$, put

- $\mathcal{O}_X^G(U) \triangleq$ the subring of all $x \in \mathcal{O}_X(U)$ such that $\sigma_\xi^\#(i_X(x)) = i_X(x)$ holds for every $\sigma \in G$.

2.5. Galois covers. Let $f : X \rightarrow Y$ be a surjective morphism of integral schemes. Note that here f is not necessarily of finite type.

Definition 2.6. X is **Galois** over Y by f if the two conditions are satisfied:

- $k(X)$ is naturally Galois over $k(Y)$ by f ;
- $\text{Aut}(X/Y) \cong \text{Gal}(k(X)/k(Y))$ are isomorphic in a natural manner.

Let $D \subseteq D_1 \cap D_2$ be three integral domains contained in a field Ω . The ring D_1 is said to be a **conjugation** of D_2 over D if there is a k -isomorphism

$$\tau : \text{Fr}(D_1) \rightarrow \text{Fr}(D_2)$$

between fields such that $\tau(D_1) = D_2$, where $k \triangleq \text{Fr}(D)$.

Definition 2.7. X is **locally quasi-galois** over Y by f if the below condition is satisfied:

For any affine open subsets $V \subseteq Y$ and $U \subseteq f^{-1}(V)$, the ring $\mathcal{O}_X(U)$ is contained in a common algebraic closure Ω of $k(X)$ and there is one and only one conjugation of $\mathcal{O}_X(U)$ over the ring $f^\#(i_Y(\mathcal{O}_Y(V)))$.

Definition 2.8. A reduced affine covering $\mathcal{C}_X$ of X with values in an algebraic closure Ω of $k(X)$ is said to be **quasi-galois closed** over Y by f if there is a local chart $(U'_\alpha, \phi'_\alpha; A'_\alpha) \in \mathcal{C}_X$ such that $U'_\alpha \subseteq \varphi^{-1}(V_\alpha)$ and $H_\alpha = A'_\alpha$ hold for

- any $(U_\alpha, \phi_\alpha; A_\alpha) \in \mathcal{C}_X$;
- any affine open set V_α in Y with $U_\alpha \subseteq f^{-1}(V_\alpha)$;
- any conjugation H_α of A_α over $B_\alpha \triangleq f^\#(i_Y(\mathcal{O}_Y(V_\alpha)))$.

Definition 2.9. X is **quasi-galois closed** (or **qc** for short) over Y by f if there is a reduced affine covering $\mathcal{C}_X$ of X with values in an algebraic closure Ω of $k(X)$ such that $\mathcal{C}_X$ is quasi-galois closed over Y by f .

Definition 2.10. X is **locally complete** over Y by f if for every automorphism $\rho \in \text{Gal}(k(X)/f^\#(k(Y)))$ and every affine open sets $V \subseteq Y$ and $U \subseteq f^{-1}(V)$, there is an affine open set $W \subseteq f^{-1}(V)$ and an isomorphism $\tilde{\rho} : \mathcal{O}_X(U) \rightarrow \mathcal{O}_X(W)$ between algebras over $f^\#(\mathcal{O}_Y(V))$ such that ρ is induced from $\tilde{\rho}$ in a natural manner.

Immediately, by definitions we have

$$\text{locally quasi-galois} \implies \text{locally complete};$$

$$\text{quasi-galois closed} \implies \text{locally complete}.$$

On the other hand, from *Step 4* in §3.3 it is seen that the above condition local completeness is automatically satisfied for integral affine schemes and for integral normal schemes.

3. A CRITERION OF GALOIS COVERS

In this section we will give a criterion of Galois covers of integral varieties, which is one of the key points for us to compute the étale fundamental group of an arithmetic variety in the paper. In §4 by such a result we will prove that there is a universal Galois cover for the étale fundamental group.

3.1. A criterion of Galois covers. The following theorem gives a criterion of Galois covers for integral varieties, which will be frequently used in the paper.

Theorem 3.1. *Let X and Y be integral varieties. Suppose that $\phi : X \rightarrow Y$ is a surjective morphism of schemes such that X is locally complete over Y and $k(X)$ is canonically an algebraic Galois extension over $k(Y)$ by ϕ . Then there is a natural isomorphism*

$$\text{Aut}(X/Y) \cong \text{Gal}(k(X)/k(Y))$$

between groups. In particular, X is a Galois cover of Y .

Remark 3.2. The conclusion of *Theorem 3.1* can be regarded as a generalization of several well-known related results in [9, 11, 16, 17, 18] for the case of integral varieties since the structure morphism ϕ here is not necessarily of finite type.

3.2. Applications of the criterion. Let X and Y be integral varieties of the same dimension. Let $f : X \rightarrow Y$ be a surjective morphism.

Lemma 3.3. *Assume that X is locally quasi-galois over Y by f . Then X is qc over Y .*

Proof. It is immediate from definition. $\square$

Lemma 3.4. *Let X be qc over Y by f . Then the function field $k(X)$ is canonically Galois over $f(Y)$; moreover, X is Galois over Y by f .*

Proof. It is immediate from *Theorem 3.1*. $\square$

Lemma 3.5. *Suppose that for any affine open set V of Y , the set $f^{-1}(V)$ is affine open in X and the ring $f^\#(i_Y(\mathcal{O}_Y(V)))$ is equal to $i_X(\mathcal{O}_X^{\text{Aut}(X/Y)}(f^{-1}(V)))$. Then X is Galois over Y by f .*

Proof. It is immediate from *Lemmas 3.4-5*. $\square$

Lemmas 3.3-5 will be used in the following section.

3.3. Proof of Theorem 3.1. Now we prove *Theorem 3.1*.

Proof. (Proof of Theorem 3.1) The process of the proof here is based on a trick originally in [1] and a refinement in [6].

Without loss of generality, by *Lemma 2.3* suppose that X and Y are essentially affine in the function fields $k(X)$ and $k(Y)$, respectively.

Here, the function field $k(X)$ will be taken as the set of elements of the form (U, f) . We will identify the function field $k(Y)$ with the subfield $\phi^\#(k(Y))$ of $k(X)$.

In the following we will proceed in several steps to prove that there is a group isomorphism

$$\text{Aut}(X/Y) \cong \text{Gal}(k(X)/k(Y)).$$

Step 1. Construct a map

$$t : \text{Aut}(X/Y) \longrightarrow \text{Gal}(k(X)/k(Y))$$

between sets.

In fact, given an automorphism

$$\sigma = (\sigma, \sigma^\#) \in \text{Aut}_k(X/Y),$$

i.e.,

$$\sigma : X \longrightarrow X$$

is a homeomorphism;

$$\sigma^\# : \mathcal{O}_X \rightarrow \sigma_* \mathcal{O}_X$$

is an isomorphism of sheaves of rings on X .

Let ξ be the generic point of X . We have

$$\sigma(\xi) = \xi;$$

it follows that

$$\sigma^\# : k(X) = \mathcal{O}_{X,\xi} \rightarrow \sigma_* \mathcal{O}_{X,\xi} = k(X)$$

is an automorphism of $k(X)$.

Denote by $\sigma^{\#-1}$ the inverse of the ring isomorphism

$$\sigma^\# : k(X) \rightarrow k(X).$$

Take any open subset U of X . We have the restriction

$$\sigma = (\sigma, \sigma^\#) : (U, \mathcal{O}_X|_U) \longrightarrow (\sigma(U), \mathcal{O}_X|_{\sigma(U)})$$

of open subschemes. That is,

$$\sigma^\# : \mathcal{O}_X|_{\sigma(U)} \rightarrow \sigma_* \mathcal{O}_X|_U$$

is an isomorphism of sheaves on $\sigma(U)$. In particular,

$$\sigma^\# : \mathcal{O}_X(\sigma(U)) = \mathcal{O}_X|_{\sigma(U)}(\sigma(U)) \rightarrow \mathcal{O}_X(U) = \sigma_* \mathcal{O}_X|_U(\sigma(U))$$

is an isomorphism of rings.

For every $f \in \mathcal{O}_X|_U(U)$, we have

$$f \in \sigma_* \mathcal{O}_X|_U(\sigma(U));$$

hence

$$\sigma^{\#-1}(f) \in \mathcal{O}_X(\sigma(U)).$$

Now we have a map

$$t : \text{Aut}_k(X/Y) \longrightarrow \text{Gal}(k(X)/k(Y))$$

between sets given by

$$\sigma = (\sigma, \sigma^\#) \longmapsto t(\sigma) = \langle \sigma, \sigma^{\#-1} \rangle$$

such that

$$\langle \sigma, \sigma^{\#-1} \rangle : (U, f) \in \mathcal{O}_X(U) \longmapsto (\sigma(U), \sigma^{\#-1}(f)) \in \mathcal{O}_X(\sigma(U))$$

is a mapping of $k(X)$ into $k(X)$.

Step 2. Prove that the map

$$t : \text{Aut}(X/Y) \longrightarrow \text{Gal}(k(X)/k(Y))$$

is a homomorphism between groups.

In deed, given any

$$\sigma = (\sigma, \sigma^\#) \in \text{Aut}(X/Y).$$

For any $(U, f), (V, g) \in k(X)$, we have

$$(U, f) \cdot (V, g) = (U \cap V, f \cdot g);$$

$$(U, f) + (V, g) = (U \cap V, f + g).$$

Then

$$\begin{aligned} \langle \sigma, \sigma^{\#-1} \rangle((U, f) \cdot (V, g)) &= \langle \sigma, \sigma^{\#-1} \rangle((U, f)) \cdot \langle \sigma, \sigma^{\#-1} \rangle((V, g)); \\ \langle \sigma, \sigma^{\#-1} \rangle((U, f) + (V, g)) &= \langle \sigma, \sigma^{\#-1} \rangle((U, f)) + \langle \sigma, \sigma^{\#-1} \rangle((V, g)). \end{aligned}$$

Hence, $\langle \sigma, \sigma^{\sharp-1} \rangle$ is an automorphism of $k(X)$.

It needs to prove that $\langle \sigma, \sigma^{\sharp-1} \rangle$ is an automorphism over $k(Y)$. Consider the given structure morphism

$$\phi = (\phi, \phi^{\sharp}) : (X, \mathcal{O}_X) \rightarrow (Y, \mathcal{O}_Y)$$

between schemes. As $\phi(\xi)$ is the generic point of Y and ξ is invariant under any automorphism $\sigma \in \text{Aut}(X/Y)$, it is seen that $\sigma^{\sharp} : \mathcal{O}_{X,\xi} \rightarrow \mathcal{O}_{X,\xi}$ is an isomorphism of algebras over the subalgebra

$$k(Y) = \phi^{\sharp}(k(Y)) = \phi^{\sharp}(\mathcal{O}_{Y,\phi(\xi)}),$$

i.e.,

$$\langle \sigma, \sigma^{\sharp-1} \rangle|_{\phi^{\sharp}(k(Y))} = \text{id}_{\phi^{\sharp}(k(Y))}.$$

Hence,

$$t(\sigma) = \langle \sigma, \sigma^{\sharp-1} \rangle \in \text{Gal}(k(X)/k(Y)).$$

This also proves that t is well-defined as a map between sets.

On the other hand, take any two automorphisms

$$\sigma = (\sigma, \sigma^{\sharp}), \delta = (\delta, \delta^{\sharp}) \in \text{Aut}(X/Y).$$

We have

$$\delta^{\sharp-1} \circ \sigma^{\sharp-1} = (\delta \circ \sigma)^{\sharp-1}$$

according to preliminary facts on morphisms of schemes (see [10, 12, 13]).

Then

$$\langle \delta, \delta^{\sharp-1} \rangle \circ \langle \sigma, \sigma^{\sharp-1} \rangle = \langle \delta \circ \sigma, \delta^{\sharp-1} \circ \sigma^{\sharp-1} \rangle.$$

Hence,

$$t : \text{Aut}(X/Y) \rightarrow \text{Gal}(k(X)/k(Y))$$

is a homomorphism of groups.

Step 3. Prove that

$$t : \text{Aut}(X/Y) \longrightarrow \text{Gal}(k(X)/k(Y))$$

is an injective homomorphism.

In deed, take any two automorphisms $\sigma, \sigma' \in \text{Aut}(X/Y)$ such that

$$t(\sigma) = t(\sigma').$$

We have

$$(\sigma(U), \sigma^{\sharp-1}(f)) = (\sigma'(U), \sigma'^{\sharp-1}(f))$$

for each element $(U, f) \in k(X)$. In particular,

$$(\sigma(U_0), \sigma^{\sharp-1}(f)) = (\sigma'(U_0), \sigma'^{\sharp-1}(f))$$

holds for any $f \in \mathcal{O}_X(U_0)$ and any affine open subset U_0 of X .

It is seen that there are three rings

$$A_0 = \mathcal{O}_X(U_0);$$

$$B_0 = \mathcal{O}_X(\sigma(U_0));$$

$$B_0' = \mathcal{O}_X(\sigma'(U_0)),$$

as subrings of $k(X)$, satisfying the property

$$B_0 = \sigma^{\sharp-1}(A_0) = \sigma'^{\sharp-1}(A_0) = B_0'.$$

Then

$$\sigma|_{U_0} = \sigma'|_{U_0}$$

holds as isomorphisms of schemes in virtue of preliminary facts on morphisms of affine schemes (see [10, 12, 13]).

Let U_0 run through all affine open sets of X . We must have

$$\sigma = \sigma'$$

on the whole of X . This proves that t is an injection.

Step 4. Prove that

$$t : \text{Aut}(X/Y) \longrightarrow \text{Gal}(k(X)/k(Y))$$

is a surjective homomorphism.

In fact, fixed an automorphism $\rho \in \text{Gal}(k(X)/k(Y))$. As

$$k(X) = \{(U_f, f) : f \in \mathcal{O}_X(U_f) \text{ and } U_f \subseteq X \text{ is open}\},$$

we have

$$\rho : (U_f, f) \in k(X) \longmapsto (U_{\rho(f)}, \rho(f)) \in k(X)$$

according to *Proposition 1.44* of [13], where U_f and $U_{\rho(f)}$ are open sets in X , f is contained in $\mathcal{O}_X(U_f)$, and $\rho(f)$ is contained in $\mathcal{O}_X(U_{\rho(f)})$.

We will proceed in several sub-steps to prove that for each element

$$\rho \in \text{Gal}(k(X)/k(Y))$$

there is a unique element

$$\lambda \in \text{Aut}(X/Y)$$

such that

$$t(\lambda) = \rho.$$

Substep 4-a. Fixed any affine open set V of Y . Show that for each affine open set $U \subseteq \phi^{-1}(V)$ there is an affine open set U_ρ in X such that ρ determines an isomorphism between affine schemes $(U, \mathcal{O}_X|_U)$ and $(U_\rho, \mathcal{O}_X|_{U_\rho})$.

In fact, take any affine open sets $V \subseteq Y$ and $U \subseteq \phi^{-1}(V)$. We have

$$A = \mathcal{O}_X(U) = \{(U_f, f) \in k(X) : U_f \supseteq U\}.$$

Put

$$B = \{(U_{\rho(f)}, \rho(f)) \in k(X) : (U_f, f) \in A\}.$$

Then there exists an affine open set $W \subseteq \phi^{-1}(V)$ and an isomorphism

$$\tilde{\rho} = \rho|_A : A \rightarrow B = \mathcal{O}_X(W)$$

such that ρ induces naturally from $\tilde{\rho}$ according to the assumption that X is locally complete over Y . Write $U_\rho \triangleq W$.

Therefore, by ρ we have a unique isomorphism

$$\lambda_U = \left(\lambda_U, \lambda_U^\sharp \right) : (U, \mathcal{O}_X|_U) \rightarrow (U_\rho, \mathcal{O}_X|_{U_\rho})$$

of the affine open subscheme in X such that

$$\rho|_{\mathcal{O}_X(U)} = \lambda_U^{\sharp-1} : \mathcal{O}_X(U) \rightarrow \mathcal{O}_X(U_\rho).$$

Substep 4-b. Take any affine open sets $V \subseteq Y$ and $U, U' \subseteq \phi^{-1}(V)$. Show that

$$\lambda_U|_{U \cap U'} = \lambda_{U'}|_{U \cap U'}$$

holds as morphisms of schemes.

In fact, by the above construction in *Substep 4-a*, for each λ_U it is seen that $\lambda_U^\sharp$ and $\lambda_{U'}^\sharp$ coincide restricted to the intersection $U \cap U'$ as homomorphisms of rings since we have

$$\begin{aligned}\rho|_{\mathcal{O}_X(U \cap U')} &= \lambda_U|_{U \cap U'}^{\sharp-1} : \mathcal{O}_X(U \cap U') \rightarrow \mathcal{O}_X((U \cap U')_\rho); \\ \rho|_{\mathcal{O}_X(U \cap U')} &= \lambda_{U'}|_{U \cap U'}^{\sharp-1} : \mathcal{O}_X(U \cap U') \rightarrow \mathcal{O}_X((U \cap U')_\rho).\end{aligned}$$

On the other hand, for any point $x \in U \cap U'$, we must have

$$\lambda_U(x) = \lambda_{U'}(x).$$

Otherwise, if $\lambda_U(x) \neq \lambda_{U'}(x)$, will have an affine open subset X_0 of X that contains either of the points $\lambda_U(x)$ and $\lambda_{U'}(x)$ but does not contain the other since the underlying space of X is a Kolmogorov space (see [10, 12, 13]). Assume $\lambda_U(x) \in X_0$ and $\lambda_{U'}(x) \notin X_0$. We choose an affine open subset U_0 of X such that $x \in U_0 \subseteq U \cap U'$ and $\lambda_U(U_0) \subseteq X_0$ since we have

$$\begin{aligned}\lambda_U(U \cap U') &= (U \cap U')_\rho \subseteq U_\rho; \\ \lambda_{U'}(U \cap U') &= (U \cap U')_\rho \subseteq U'_\rho.\end{aligned}$$

However, for each λ_U , we have

$$\lambda_U(U_0) = (U_0)_\rho = \lambda_{U'}(U_0);$$

then

$$\lambda_{U'}(x) \in (U_0)_\rho \subseteq X_0,$$

where there will be a contradiction.

Hence, λ_U and $\lambda_{U'}$ coincide on $U \cap U'$ as mappings of topological spaces.

Substep 4-c. By gluing the λ_U along all such affine open subsets U , we have a homeomorphism λ of X onto X as a topological space given by

$$\lambda : x \in X \mapsto \lambda_U(x) \in X$$

where x belongs to U and U is an affine open subset of X . We have

$$\lambda|_U = \lambda_U.$$

By *Substep 4-b* it is seen that λ is well-defined. Then we have an automorphism, namely λ , of the scheme $(X, \mathcal{O}_X)$ (see [10, 12, 13]).

Substep 4-d. Show that λ is contained in $\text{Aut}(X/Y)$ satisfying

$$t(\lambda) = \rho.$$

In deed, as ρ is an isomorphism of $k(X)$ over $k(Y)$, the isomorphism λ_U is over Y by ϕ for any affine open subset U of X ; then λ is an automorphism of X over Y by ϕ such that

$$t(\lambda) = \rho.$$

This proves that for each $\rho \in \text{Gal}(k(X)/k(Y))$, there exists $\lambda \in \text{Aut}(X/Y)$ such that

$$t(\lambda) = \rho.$$

Hence, t is surjective.

Therefore, by *Steps 1-4* above, it is seen that there is a group isomorphism

$$t : \text{Aut}(X/Y) \longrightarrow \text{Gal}(k(X)/k(Y))$$

by

$$\sigma = (\sigma, \sigma^\sharp) \longmapsto t(\sigma) = \langle \sigma, \sigma^{\sharp-1} \rangle,$$

where $\langle \sigma, \sigma^{\sharp-1} \rangle$ is the map of $k(X)$ into $k(X)$ given by

$$(U, f) \in \mathcal{O}_X(U) \subseteq k(X) \mapsto (\sigma(U), \sigma^{\sharp-1}(f)) \in \mathcal{O}_X(\sigma(U)) \subseteq k(X)$$

for any open set U in X and any element $f \in \mathcal{O}_X(U)$.

This completes the proof. $\square$

4. THE CALCULUS AND CONSTRUCTION OF GALOIS COVERS

In this section we will give a universal construction of Galois covers over a given integral variety. For an arithmetic variety X , in general, we will have many distinct Galois covers X_{et} over X for the étale fundamental group $\pi_1^{et}(X; s)$ such that each X_{et} is a realization for the group $\pi_1^{et}(X; s)$, i.e., there will be isomorphisms

$$\begin{aligned} k(X_{et}) &\cong k(X; \Omega)^{un}; \\ Gal(k(X; \Omega)^{un}/k(X)) &\cong Aut(X_{et}/X) \end{aligned}$$

both in a natural manner.

Furthermore, all these Galois covers X_{et} of X can have the same function fields $k(X; \Omega)^{un}$ but can not be isomorphic as schemes. In other words, for any such Galois covers X_{et} and X'_{et} over X , we have

$$\begin{aligned} X_{et} &\not\cong X'_{et}; \\ k(X_{et}) &\cong k(X'_{et}); \\ Aut(X_{et}/X) &\cong Aut(X'_{et}/X). \end{aligned}$$

4.1. Calculus and existence of Galois covers. Here we give the existence of Galois covers of an integral variety such that the Galois covers can be prescribed by any given Galois extensions of the function field.

Theorem 4.1. *For a field L and an integral variety Y such that L is an algebraic Galois extension of $k(Y)$, there exists an integral variety X_L and an affine surjective morphism $f_L : X_L \rightarrow Y$ satisfying the properties:*

- $L = k(X_L)$;
- X_L is essentially affine in L ;
- X_L is Galois over Y by f_L ;
- X_L is locally quasi-galois over Y by f_L ;
- $Aut(X_L/Y)$ acts on the fiber $f_L^{-1}(y)$ transitively for any $y \in Y$;
- For any affine open set V of Y , there is an isomorphism

$$\mathcal{O}_Y(V) \cong \mathcal{O}_{X_L}^{Aut(X_L/Y)}(f_L^{-1}(V))$$

of groups in a natural manner.

The above integral variety X_L with a morphism f_L , denoted by (X_L, f_L) , will be called a **Galois cover of Y in a field L** .

Now consider the calculus of Galois covers of an integral variety.

Definition 4.2. Let Y be an integral variety and let L be an algebraic Galois extension of $k(Y)$. Put

- $\Sigma[L/Y] \triangleq$ the collection of all the sets Δ of generators of L over $k(Y)$ such that $\Delta \subseteq L \setminus k(Y)$;
- $Y[\Delta] \triangleq$ the integral variety X , a Galois cover of Y in L , obtained by adding a set $\Delta = \Delta(L/k(Y)) \in \Sigma[L/Y]$ to an essentially affine realization of Y in the universal construction of Galois covers in §4.2 below.

It is clear that $Y[\Delta]$ has all the properties in *Theorem 4.1* above.

Remark 4.3. Fixed an integral variety Y and a Galois extension L of $k(Y)$. There are the following statements.

- Different sets $\Delta \in \Sigma[L/Y]$ can produce different Galois covers $Y[\Delta]$ of Y in L . In general, it is not true that $Y[\Delta]$ and $Y[\Delta']$ are isomorphic over Y for any $\Delta, \Delta' \in \Sigma[L/Y]$.
- There is a biggest Galois cover $Y[\Delta_{min}]$ of Y in L , called **sp-complete**, which contains all possible points in the underlying space. For instance, take a minimal element Δ_{min} of $\Sigma[L/Y]$.
- There is a smallest Galois cover $Y[\Delta_{max}]$ of Y in L , which has the same underlying space as Y . For instance, take a maximal element Δ_{max} of $\Sigma[L/Y]$.

Proposition 4.4. Let Z be an integral variety and L an algebraic Galois extension of $k(Z)$. Then for any $\Delta, \Delta' \in \Sigma[L/Z]$, there is an isomorphism

$$Z[\Delta, \Delta^{-1}] \cong Z[\Delta', \Delta'^{-1}]$$

between schemes over Z . In particular, each $Z[\Delta, \Delta^{-1}]$ is a smallest Galois cover of Z . Here, $Z[\Delta, \Delta^{-1}]$ denotes $Z[\Delta \cup \Delta^{-1}]$.

Proof. It reduces to consider the affine scheme $Z = \text{Spec}(A)$ according to the construction in §4.2 below. But this is trivial. $\square$

4.2. A universal construction of Galois covers. In the following we will prove *Theorem 4.1*.

Proof. (Proof of Theorem 4.1) Here we take the trick in [4].

From *Lemma 2.3*, assume that Y is essentially affine in the function field $k(Y)$ without loss of generality.

Denote by $\mathcal{C}_Y$ the set of local charts (V, ψ_V, B_V) such that $B_V = \mathcal{O}_Y(V)$, where V runs through all affine open sets of Y . Let

$$\Delta(L/k(Y)) \subseteq L \setminus k(Y)$$

be a set of generators of the field L over the function field $k(Y)$.

Following the procedure developed in [4], we will proceed in several steps to construct an integral variety $X = X_L$ and a morphism $f = f_L : X_L \rightarrow Y$.

Step 1. For any local chart $(V, \psi_V, B_V) \in \mathcal{C}_Y$, define

$$A_V \triangleq B_V[\Delta_V],$$

i.e., the subring of L generated over B_V by the set

$$\Delta_V \triangleq \{\sigma(w) \in L : \sigma \in \text{Gal}(L/k(Y)), w \in \Delta(L/k(Y))\}.$$

Then $\text{Fr}(A_V) = L$ holds. It is seen that B_V is exactly the invariant subring of the natural action of the Galois group $\text{Gal}(L/M)$ on A_V .

Set

$$i_V : B_V \rightarrow A_V$$

to be the inclusion.

Step 2. Define the disjoint union

$$\Sigma = \coprod_{(V, \psi_V, B_V) \in \mathcal{C}_Y} \text{Spec}(A_V).$$

Then Σ is a topological space, where the topology τ_Σ on Σ is naturally determined by the Zariski topologies on all $\text{Spec}(A_V)$.

Let

$$\pi_Y : \Sigma \rightarrow Y$$

be the projection.

Step 3. Given an equivalence relation R_Σ in Σ in such a manner:

For any $x_1, x_2 \in \Sigma$, we say

$$x_1 \sim x_2$$

if and only if the two conditions are satisfied:

- $\pi_Y(x_1) = \pi_Y(x_2)$ holds as points in Y ;
- x_1 and x_2 are L -equivalent as affine L -points.

Let

$$X = \Sigma / \sim$$

and let

$$\pi_X : \Sigma \rightarrow X$$

be the projection.

It is seen that X is a topological space as a quotient of Σ .

Step 4. Define a map

$$f : X \rightarrow Y$$

by

$$\pi_X(z) \longmapsto \pi_Y(z)$$

for each $z \in \Sigma$.

Step 5. Define

$$\mathcal{C}_X = \{(U_V, \varphi_V, A_V)\}_{(V, \psi_V, B_V) \in \mathcal{C}_Y}$$

where $U_V \triangleq \pi_Y^{-1}(V)$ is an open set in X and $\varphi_V : U_V \rightarrow \text{Spec}(A_V)$ is the identity map on U_V for each $(V, \psi_V, B_V) \in \mathcal{C}_Y$.

We have an integral scheme $(X, \mathcal{O}_X)$ by gluing the affine schemes $\text{Spec}(A_V)$ for all local charts $(V, \psi_V, B_V) \in \mathcal{C}_Y$ with respect to the equivalence relation R_Σ (see [10, 12]).

It is seen that $\mathcal{C}_X$ is a reduced affine covering of X with values in L . In particular, $\mathcal{C}_X$ is quasi-galois closed over Y by f .

By *Lemma 3.5* it is seen that X and f have the desired properties.

This completes the proof. $\square$

4.3. A realization of the étale fundamental group. From *Theorem 4.1* we have the following realization for the étale fundamental group of an arithmetic scheme.

Lemma 4.5. *Let X be an arithmetic variety and Ω an algebraic closure of $k(X)$. Then there exists an integral variety X_{et} and an affine surjective morphism*

$$f_{et} : X_{et} \rightarrow X$$

such that

- $k(X_{et}) = k(X; \Omega)^{un}$;
- (X_{et}, f_{et}) is a Galois cover of X in the sense of *Theorem 4.1*.

Such a Galois cover (X_{et}, f_{et}) will be called a **universal Galois cover** of X for the étale fundamental group $\pi_1^{et}(X, s)$. Here s is a geometric point of X over Ω .

Remark 4.6. Let X be an arithmetic variety and Ω an algebraic closure of $k(X)$. Then for any

$$\Delta_{un} \in \Sigma[k(X; \Omega)^{un}/X],$$

$X[\Delta_{un}]$ is a universal Galois cover of X for the étale fundamental group $\pi_1^{et}(X, s)$. In general, there can be many distinct universal Galois covers of X for $\pi_1^{et}(X, s)$ according to the calculus of Galois covers in §4.1.

5. FORMALLY UNRAMIFIED = ARITHMETICALLY UNRAMIFIED

In this section we will prove that for an arithmetic scheme, the maximal formally unramified extension is equal to the maximal arithmetically unramified extension.

At the same time, arithmetically unramified, which is defined in an evident manner, coincides with that in algebraic number theory and has many arithmetic operations and properties. Hence, via arithmetically unramified, it is seen the maximal formally unramified extension is exactly of an arithmetic sense.

On the other hand, it is seen that arithmetically unramified is more accessible to operations (see [7, 8]); it follows that arithmetically unramified extensions can serve as a computation of the étale fundamental group of an arithmetic variety.

5.1. Arithmetic function fields and arithmetically unramified extensions. Recall that an **arithmetic function field over variables** $t_1, t_2, \dots, t_n$ is the fractional field of an integral domain

$$\mathbb{Z}[t_1, t_2, \dots, t_n, s_1, s_2, \dots, s_m]$$

of finite type over $\mathbb{Z}$, where

$$s_1, s_2, \dots, s_m$$

are algebraic over $\mathbb{Q}[t_1, t_2, \dots, t_n]$ and

$$t_1, t_2, \dots, t_n,$$

are a finite number of (algebraic independent) variables over $\mathbb{Q}$.

Definition 5.1. (see [5, 7, 8]) Let K be an arithmetic function field over variables $t_1, t_2, \dots, t_n$. As in algebraic number theory, put

- $\mathcal{O}_{[t_1, t_2, \dots, t_n]K} \triangleq$ the subring of elements $x \in K$ that are integral over the integral domain $\mathbb{Z}[t_1, t_2, \dots, t_n]$.

The ring $\mathcal{O}_{[t_1, t_2, \dots, t_n]K}$ is said to be the **ring of algebraic integers of K over variables** $t_1, t_2, \dots, t_n$.

It is easily seen that $K = Fr(\mathcal{O}_K)$ holds. Sometimes, we will write

$$\mathcal{O}_K = \mathcal{O}_{[t_1, t_2, \dots, t_n]K}$$

for brevity.

Let K and L be two arithmetic function fields over variables $t_1, t_2, \dots, t_n$. We say

$$(L, \mathcal{O}_L) \supseteq (K, \mathcal{O}_K)$$

if and only if there are the properties

$$\overline{K} \supseteq L \supseteq K;$$

$$\mathcal{O}_L \supseteq \mathcal{O}_K \supseteq \mathbb{Z}[t_1, t_2, \dots, t_n].$$

Definition 5.2. (see [5, 7, 8]) Let $L \supseteq K$ be any two arithmetic function fields over variables $t_1, t_2, \dots, t_n$. The field L is **arithmetically unramified** over K (**relative to variables** $t_1, t_2, \dots, t_n$) if for the pairs $(L, \mathcal{O}_L) \supseteq (K, \mathcal{O}_K)$, the affine scheme $\text{Spec}(\mathcal{O}_L)$ is unramified over $\text{Spec}(\mathcal{O}_K)$ by the morphism induced from the inclusion map $K \hookrightarrow L$.

It is clear that the arithmetically unramified extensions coincide exactly with those in algebraic number theory (for instance, see [14]).

Lemma 5.3. (see [5, 7, 8]) *The arithmetically unramified extension is independent of the choice of variables $t_1, t_2, \dots, t_n$.*

Proof. It is immediate from operations on arithmetically unramified extensions in §5.2 below. $\square$

5.2. Operations on arithmetically unramified extensions. As in algebraic number theory, we have several operations on arithmetically unramified extensions such as base changes, composites, subfields, and transitivity.

Let $K \subseteq L \subseteq M$ be arithmetic function fields over variables $t_1, t_2, \dots, t_n$. We have the following lemmas from preliminary facts on unramified morphisms and étale morphisms between schemes (see [9, 11, 15, 18]).

Lemma 5.4. (see [5, 7, 8]) *Let L/K and M/L be arithmetically unramified. Then so is M/K .*

Lemma 5.5. (see [5, 7, 8]) *Let L be arithmetically unramified over K . Take a subfield $L_0 \subseteq L$. Then L/L_0 and L_0/K are arithmetically unramified.*

Lemma 5.6. (see [5, 7, 8]) *Let L_1 and L_2 be two arithmetic function fields over variables $t_1, t_2, \dots, t_n$. Suppose that L_1 and L_2 are arithmetically unramified over K , respectively. Then the composite $L_1 \cdot L_2$ of fields is also arithmetically unramified over K .*

Lemma 5.7. (see [5, 7, 8]) *Let $K' \supseteq K$ be an arithmetic function field over variables $t_1, t_2, \dots, t_n$. Suppose that L is arithmetically unramified over K . Then the composite $L' = L \cdot K'$ of fields is also arithmetically unramified over K' .*

5.3. The maximal formally unramified extension is nothing other than the maximal arithmetically unramified extension. Now take the maximal arithmetically unramified extension of an arithmetic variety as one does in algebraic number theory, which will be proved to be equal to the maximal formally unramified extension of the arithmetic variety.

Let L be an arithmetic function field over variables $t_1, t_2, \dots, t_n$. Let Ω be an algebraic closure of L .

Put

- $[L](\Omega)^{au} \triangleq$ the set of all finite arithmetically unramified subextensions of L contained in Ω .
- $(L; \Omega)^{au} \triangleq \lim_{\rightarrow Z \in [L](\Omega)^{au}} \lambda_Z(Z)$, i.e., the direct limit of the direct system of rings indexed by $[L](\Omega)^{au}$, where each $\lambda_Z : Z \in [L](\Omega)^{au} \rightarrow \Omega$ is the L -embedding,

where $[L; \Omega]^{au}$ is taken as a directed set defined by set inclusion.

The subfield $(L; \Omega)^{au}$ of Ω is called the **maximal arithmetically unramified extension** of L in Ω .

Write

- $k(X; \Omega)^{au} \triangleq (k(X); \Omega)^{au}$ for an arithmetic variety X and an algebraic closure Ω of the function field $k(X)$.

Lemma 5.8. (see [5, 7, 8]) *Let $[L](\Omega)_0^{au}$ be the set of all finite arithmetically unramified Galois subextensions of L contained in Ω . Then we have*

$$(L; \Omega)^{au} = \lim_{\rightarrow Z \in [L](\Omega)_0^{au}} \lambda_Z(Z).$$

Moreover, $(L; \Omega)^{au}$ is an algebraic Galois extension of L .

Proof. It is immediate from the preliminary fact that $[L](\Omega)_0^{au}$ is a cofinal directed subset of $[L](\Omega)^{au}$. $\square$

Fortunately there exists the following lemma which says that for an arithmetic variety, the maximal arithmetically unramified extension is equal to the maximal formally ramified extension.

Lemma 5.9. *Let X be an arithmetic variety and let Ω be an algebraic closure of the function field $k(X)$. Then we have*

$$k(X; \Omega)^{au} = k(X; \Omega)^{un}$$

as subfields of Ω .

Proof. Let $L = k(X)$. As L is an arithmetic function field over several variables, consider the ring $\mathcal{O}_L$ of algebraic integers of L . We have

$$L = k(\text{Spec}(\mathcal{O}_L));$$

then

$$(L; \Omega)^{un} = (k(\text{Spec}(\mathcal{O}_L)); \Omega)^{un}.$$

To prove the lemma, it suffices to prove

$$(k(\text{Spec}(\mathcal{O}_L)); \Omega)^{un} = (L; \Omega)^{au}$$

holds.

In deed, we have

$$\begin{aligned} (L; \Omega)^{au} &= \lim_{\rightarrow Z \in [L](\Omega)_0^{au}} \lambda_Z(Z); \\ (k(\text{Spec}(\mathcal{O}_L)); \Omega)^{un} &= \lim_{\rightarrow X \in [\text{Spec}(\mathcal{O}_L); \Omega]_{et}} \lambda_X(k(X)). \end{aligned}$$

On the other hand, put

$$\Sigma = \{k(X) \mid X \in [\text{Spec}(\mathcal{O}_L); \Omega]_{et}\}.$$

Then Σ is a directed set, where the partial order is given by set inclusion.

As $\text{Spec}(\mathcal{O}_L)$ is a normal scheme, it is easily seen that each $X \in [\text{Spec}(\mathcal{O}_L); \Omega]_{et}$ is normal from preliminary facts on étale covers (see [11, 15, 18]). It follows that $[L](\Omega)_0^{au}$ is a cofinal directed subset of Σ .

Hence,

$$\begin{aligned} &(k(\text{Spec}(\mathcal{O}_L)); \Omega)^{un} \\ &= \lim_{\rightarrow X \in [\text{Spec}(\mathcal{O}_L); \Omega]_{et}} \lambda_X(k(X)) \\ &= \lim_{\rightarrow Z \in \Sigma} \lambda_Z(Z) \\ &= \lim_{\rightarrow Z \in [L](\Omega)_0^{au}} \lambda_Z(Z) \\ &= (L; \Omega)^{au}. \end{aligned}$$

This completes the proof. $\square$

By Lemma 5.9 we have such a computation of the étale fundamental group of an arithmetic variety.

Remark 5.10. For any arithmetic variety X , there is an isomorphism

$$\pi_1^{et}(X, s) \cong \text{Gal}(k(X; \Omega)^{au}/k(X))$$

of groups in a natural manner. Here, s is a geometric point of X over an algebraic closure Ω of the function field $k(X)$.

6. PROOF OF THE MAIN THEOREM

6.1. Recalling the étale fundamental group. Fixed a connected scheme Z . By a **finite étale cover** of Z we understand a scheme that is finite and étale over Z by a surjective morphism. Denote by

$$\text{Fet}[Z]$$

the set of all finite étale covers of Z .

Fixed a geometric point s of Z over a given separable closure Ω of $k(Z)$. There is a fiber functor Fib_s on $\text{Fet}[Z]$ given by

$$\text{Fib}_s(Y) = \text{sp}(Y \times_Z \text{Spec}(\Omega))$$

for any $Y \in \text{Fet}[Z]$. Here $\text{sp}(W)$ denotes the underlying space of a scheme W .

Then the **étale fundamental group** of Z over s , denoted by $\pi_1^{et}(Z; s)$, is defined to be the group of automorphisms of the fiber functor Fib_s over Z . See [11, 15, 18] for details.

Let Y be a finite étale cover of Z by a morphism f and let t be a geometric point of Y over s with $s = f \circ t$. Then (Y, t) is said to be a **pointed finite étale cover** of (Z, s) .

Remark 6.1. Let $\text{Fet}[Z; \Omega]$ denote the set of pointed finite étale covers of (Z, s) . For any $(Y_1, t_1), (Y_2, t_2) \in \text{Fet}[Z; \Omega]$, we say $(Y_1, t_1) \leq (Y_2, t_2)$ if and only if (Y_2, t_2) is a pointed finite étale cover of (Y_1, t_1) . Then $\text{Fet}[Z; \Omega]$ is a directed set with $\leq$.

Remark 6.2. There is an isomorphism

$$\pi_1^{et}(Z; s) \cong \lim_{\leftarrow (Y, t) \in \text{Fet}[Z; \Omega]} \text{Aut}(Y/Z)$$

as inverse limits of inverse systems of groups.

In practice, we usually use finite étale Galois covers to compute the inverse limit, i.e., the étale fundamental group. Recall that $(Y, t) \in \text{Fet}[Z; \Omega]$ is said to be a **finite étale Galois cover** of (Z, s) if Y is connected and $\#\text{Aut}(Y/Z)$ is equal to $\#\text{Fib}_s(Y)$. It is seen that $\#\text{Aut}(Y/Z)$ and $\#\text{Fib}_s(Y)$ are equal if and only if $\text{Aut}(Y/Z)$ induces a transitive action on $\text{Fib}_s(Y)$ (see [11, 15, 18]).

Let $[X; \Omega]_{et}^*$ be the set of finite étale Galois covers of (Z, s) , where any two $(Y_1, t_1), (Y_2, t_2) \in [X; \Omega]_{et}^*$ are identified with each other if there is an isomorphism of (Y_1, t_1) onto (Y_2, t_2) over (Z, s) .

Remark 6.3. As a directed set, $[X; \Omega]_{et}^*$ is isomorphic to a cofinal subset of $\text{Fet}[Z; \Omega]$.

Remark 6.4. There are isomorphisms

$$\pi_1^{et}(Z; s) \cong \lim_{\leftarrow (Y, t) \in \text{Fet}[Z; \Omega]} \text{Aut}(Y/Z) \cong \lim_{\leftarrow (Y, t) \in [X; \Omega]_{et}^*} \text{Aut}(Y/Z)$$

as inverse limits of inverse systems of groups.

6.2. A preliminary fact. Now let X be an arithmetic variety. It is seen that X is connected since an irreducible space must be connected. Let s be a geometric point of X over a given separable closure Ω of $k(X)$. As introduced in §1, $[X; \Omega]_{et}$ consists of elements $(Y, y) \in [X; \Omega]_{et}^*$ such that Y is irreducible. Then we have

$$[X; \Omega]_{et} \subseteq [X; \Omega]_{et}^*$$

as a subset.

Lemma 6.5. *Every finite étale Galois cover of an arithmetic variety is irreducible. That is, for any arithmetic variety X , we have*

$$[X; \Omega]_{et}^* = [X; \Omega]_{et}.$$

Proof. Let $(Y, y) \in [X; \Omega]_{et}^*$. By taking a section of Y over X , it is seen that the underling spaces

$$sp(X) \cong sp(Y)$$

are homeomorphic since X and Y are connected. As X is irreducible, Y must be irreducible. $\square$

6.3. Proof of the main theorem. Now we can prove *Theorem 1.1*.

Proof. (Proof of Theorem 1.1) Let Ω be an algebraic closure of the function field $k(X)$. By *Lemma 2.3*, assume that X is essentially affine in Ω without loss of generality. For brevity, write

$$k(X)^{un} = k(X; \Omega)^{un}.$$

Let $\Delta \subseteq k(X)^{un} \setminus k(X)$ be a set of generators of the field $k(X)^{un}$ over $k(X)$. Put

$$\begin{aligned} I &= \{\text{finite subsets of } \Delta\}; \\ G &= \text{Gal}(k(X)^{un}/k(X)). \end{aligned}$$

We will proceed in several steps in the following.

Step 1. As in §4.1, for Δ and G we have an integral variety

$$X_{\infty}^{un} = X[\Delta]$$

and a surjective morphism

$$f_{\infty} : X_{\infty}^{un} \rightarrow X$$

such that

- $k(X_{\infty}^{un}) = k(X)^{un}$;
- X_{∞}^{un} is Galois over X by f_{∞} ;
- X_{∞}^{un} is essentially affine in $k(X; s)^{un} \subseteq \Omega$.

Step 2. Let $\alpha \in I$. Likewise, for α and $G_{\alpha} = \text{Gal}(k(X)[\alpha]/k(X))$ we have an integral variety

$$X_{\alpha}^{un} = X[\alpha]$$

and a surjective morphism

$$f_{\alpha} : X_{\alpha}^{un} \rightarrow X$$

satisfying the properties:

- $k(X_{\alpha}^{un}) \subseteq k(X)^{un}$;
- X_{α}^{un} is Galois over X by f_{α} ;
- X_{α}^{un} is essentially affine in $k(X_{\alpha}^{un})$.

Step 3. Fixed any $\alpha \subseteq \beta$ in I . We have integral varieties X_α^{un} and X_β^{un} and a surjective morphism

$$f_\alpha^\beta : X_\beta^{un} \rightarrow X_\alpha^{un}$$

of finite type, where f_α^β is obtained as in §4.2 in a natural manner and satisfies the property

$$f_\beta = f_\alpha \circ f_\alpha^\beta.$$

Then we have Galois covers

$$\begin{aligned} X_\alpha^{un}/X; \\ X_\beta^{un}/X; \\ X_\beta^{un}/X_\alpha^{un}. \end{aligned}$$

As Δ is an infinite set, there is a γ in I such that

$$\gamma \supseteq \alpha, \gamma \supseteq \beta.$$

In particular,

$$X_\gamma^{un}/X_\beta^{un}$$

and

$$X_\gamma^{un}/X_\alpha^{un}$$

are both Galois covers.

Step 4. For any $\alpha, \beta \in I$, we say $\alpha \leq \beta$ if and only if $\alpha \subseteq \beta$. Then with $\leq$, I is a partially ordered set.

We have a direct system

$$\{k(X_\alpha^{un}); i_\alpha^\beta\}_{\alpha \in I}$$

of rings, where each

$$i_\alpha^\beta : k(X_\alpha^{un}) \rightarrow k(X_\beta^{un})$$

is a homomorphism of fields induced by the morphism f_α^β .

For the fields, we have

$$k(X_\infty^{un}) = \lim_{\rightarrow \alpha \in I} k(X_\alpha^{un}).$$

For the Galois groups, we have

$$\text{Gal}(k(X_\infty^{un})/k(X)) \cong \lim_{\leftarrow \alpha \in I} \text{Gal}(k(X_\alpha^{un})/k(X)).$$

Step 5. Put

$$[X; \Omega]_{qc} = \{X_\alpha^{un} : \alpha \in I\}.$$

For any $X_\alpha^{un}, X_\beta^{un} \in [X; \Omega]_{qc}$, we say

$$X_\alpha^{un} \leq X_\beta^{un}$$

if and only if X_β^{un} is Galois over X_α^{un} .

Then $[X; \Omega]_{qc}$ is a directed set, where two elements $Y, Z \in [X; \Omega]_{qc}$ are identified with each other if they are isomorphic over X as schemes.

Consider $[X; \Omega]_{et}$, the set of finite étale Galois covers of X with geometric points over s_ξ , where two elements $Y, Z \in [X; \Omega]_{et}$ are identified with each other if they are isomorphic over X as schemes.

Without loss of generality, in the remainder of the paper, every $X_\alpha \in [X; \Omega]_{et}$ is assumed to be essentially affine in the function field $k(X_\alpha)$ from *Lemma 2.3*.

For any $X_\alpha, X_\beta \in [X; \Omega]_{et}$, we say

$$X_\alpha \leq X_\beta$$

if and only if X_β is a finite étale Galois cover over X_α . It follows that $[X; \Omega]_{et}$ is a directed set.

It is easily seen that $[X; \Omega]_{et}$ is a directed subset of $[X; \Omega]_{qc}$.

Step 6. Prove $[X; \Omega]_{et}$ is cofinal in $[X; \Omega]_{qc}$.

In deed, according to *Step 5*, we have two inverse systems of groups

$$\{Aut(Y/X) : Y \in [X; \Omega]_{et}\};$$

$$\{Aut(Z/X) : Z \in [X; \Omega]_{qc}\}.$$

Denote by

$$\eta : [X; \Omega]_{et} \hookrightarrow [X; \Omega]_{qc}$$

the inclusion map of the sets. Then η induces a map

$$\eta_* : \{Aut(Y/X) : Y \in [X; \Omega]_{et}\} \rightarrow \{Aut(Z/X) : Z \in [X; \Omega]_{qc}\}$$

of inverse systems of groups, given by

$$Aut(Y/X) \mapsto Aut(Y/X).$$

It is clear that η_* is injective.

On the other hand, take any $Z \in [X; \Omega]_{qc}$. It is seen that $k(Z)$ is a finite Galois extension of $k(X)$. As a subfield of $k(X)^{un}$, $k(Z)$ must be or contained in a finite étale Galois extension $k(Z')$ of $k(X)$ with $Z' \in [X; \Omega]_{et}$ by *Lemma 5.9* and according to the operations on arithmetically unramified extensions in §5.2.

For Z' , we have two cases:

Case (i). Let $k(Z) = k(Z')$. Then we have

$$Aut(Z'/X) = Aut(Z/X).$$

In such a case, we put

$$Z^* = Z'.$$

Case (ii). Let $k(Z) \subsetneq k(Z')$. Then $k(Z')$ is a Galois extension of $k(Z)$ and there is a $Z^* \in [X; \Omega]_{qc}$ such that

$$\begin{aligned} Z^* &\geq Z; \\ k(Z^*) &= k(Z'); \\ Aut(Z^*/X) &\cong Gal(k(Z')/k(X)) \cong Aut(Z'/X). \end{aligned}$$

Now define

$$[X; \Omega]_{qc}^* = \{Z^* : Z \in [X; \Omega]_{qc}\}$$

where each Z^* is given in the manner of the two cases above.

As a directed subset, we have

$$[X; \Omega]_{qc}^* \subseteq [X; \Omega]_{qc}.$$

It is seen that $[X; \Omega]_{et}$ is cofinal in $[X; \Omega]_{qc}^*$ and $[X; \Omega]_{qc}^*$ is cofinal in $[X; \Omega]_{qc}$. Hence, $[X; \Omega]_{et}$ is cofinal in $[X; \Omega]_{qc}$.

Step 7. As the inverse limits are isomorphic for inverse systems of groups indexed by cofinal directed sets, we have

$$\begin{aligned}
& \pi_1^{et}(X, s) \\
&= \lim_{\leftarrow} \text{Aut}(Z/X)_{Z \in [X; \Omega]_{et}} \\
&\cong \lim_{\leftarrow} \text{Aut}(Z/X)_{Z \in [X; \Omega]_{qc}} \\
&\cong \lim_{\leftarrow} \text{Gal}(k(Z)/k(X))_{Z \in [X; \Omega]_{qc}} \\
&= \lim_{\leftarrow} \text{Gal}(k(X_\alpha^{un})/k(X))_{\alpha \in I} \\
&\cong \text{Gal}(k(X_\infty^{un})/k(X)) \\
&= \text{Gal}(k(X; \Omega)^{un}/k(X)).
\end{aligned}$$

This completes the proof. $\square$

Acknowledgment. The author would like to express his sincere gratitude to Professor Li Banghe for his advice and instructions on algebraic geometry and topology.

The author would also like to express his sincere gratitude to Professor Laurent Moret-Bailly for the stimulating discussion on an earlier version of arithmetically unramified.

The present paper is a revised version of the preprint [2].

REFERENCES

- [1] An, F-W. Automorphism groups of quasi-galois closed arithmetic schemes. eprint arXiv:0907.0842.
- [2] An, F-W. On the étale fundamental groups of arithmetic schemes. eprint arXiv:0910.4646.
- [3] An, F-W. On the algebraic fundamental groups. eprint arXiv:0910.4691.
- [4] An, F-W. On the existence of geometric models for function fields in several variables. eprint arXiv:0909.1993.
- [5] An, F-W. A computation of étale fundamental groups of arithmetic schemes: special cases. preprint.
- [6] An, F-W. On the quasi-galois cover of an integral scheme. preprint.
- [7] An, F-W. On the unramified extension of an arithmetic function field in several variables. eprint arXiv:1006.5143.
- [8] An, F-W. On the unramified class fields of an arithmetic function field in several variables. preprint.
- [9] Freitag, E; Kiehl, R. Étale Cohomology and the Weil Conjecture. Springer, Berlin, 1988.
- [10] Grothendieck, A; Dieudonné, J. Éléments de Géométrie Algébrique. vols I-IV, Pub. Math. de l'IHES, 1960-1967.
- [11] Grothendieck, A; Raynaud, M. Revêtements Étales et Groupe Fondamental (SGA1). Springer, New York, 1971.
- [12] Hartshorne, R. Algebraic Geometry. Springer, New York, 1977.
- [13] Iitaka, S. Algebraic Geometry. Springer, New York, 1982.
- [14] Neukirch, J. Algebraic Number Theory. Springer, Berlin, 1999.
- [15] Milne, J. Étale Cohomology. Princeton University Press, Princeton, New Jersey, 1980.
- [16] Suslin, A; Voevodsky, V. Singular homology of abstract algebraic varieties. Invent. Math. 123 (1996), 61-94.
- [17] Suslin, A; Voevodsky, V. Relative cycles and Chow sheaves, in *Cycles, Transfers, and Motivic Homology Theories*, edited by Voevodsky, V; Suslin, A; Friedlander, E M. Annals of Math Studies, Vol 143. Princeton University Press, Princeton, NJ, 2000.
- [18] Szamuely, T. Galois Groups and Fundamental Groups. Cambridge University Press, NY, 2009.

SCHOOL OF MATHEMATICS AND STATISTICS, WUHAN UNIVERSITY, WUHAN, HUBEI 430072, PEOPLE'S REPUBLIC OF CHINA

E-mail address: fwan@amss.ac.cn